

Quantum Integrable Systems and Special Functions ¹

A.M.Perelomov ²

Institute of Theoretical
and Experimental Physics
B.Chermushkinskaya 25
117259 Moscow, Russia

Abstract

The class of quantum integrable systems associated with root systems was introduced in [OP 1977] as a generalization of the Calogero–Sutherland systems [Ca 1971], [Su 1972]. For the potential $v(q) = \kappa(\kappa - 1) \sin^{-2} q$, the wave functions of such systems are related to polynomials in l variables (l is a rank of root system) and they are a generalization of Gegenbauer polynomials and Jack polynomials [Ja 1970]. In [Pe 1998a], it was proved that the series for the product of two such polynomials is a κ -deformation of the Clebsch–Gordan series. This yields recurrence relations for these polynomials, in particular, for generalized zonal polynomials on symmetric spaces.

The present paper follows papers [Pe 1998a], [PRZ 1998]. In last of them, the recurrence relations were used to compute the explicit expressions for A_2 type polynomials, i.e., for the wave functions of the three-body Calogero–Sutherland system.

As it was shown by Ragoucy, Zaugg and the author of this paper (see [Pe 1999] and Appendix B), the similar results are also valid in A_2 case for the more general two-parameter deformation ((q, t) -deformation) introduced by Macdonald [Ma 1988].

¹Published in: Proc. Clausthal Conference 1999, *Lie Theory and Its Applications in Physics III*, pp.139-154, World Scientific, Singapore, 2000

² Current e-mail address: perelomo@dftuz.unizar.es.

1 Introduction

The class of quantum integrable systems associated with root systems was introduced in [OP 1977] (see also [OP 1978] and [OP 1983]) as a generalization of the Calogero–Sutherland systems [Ca 1971], [Su 1972]. Such systems depend on one real parameter κ (for root systems of the type A_n , D_n and E_6 , E_7 , E_8), on two parameters (for the type B_n , C_n , F_4 and G_2) and on three parameters for the type BC_n . These parameters are related to the coupling constants of the quantum system.

For the potential $v(q) = \kappa(\kappa - 1) \sin^{-2} q$ and special values of parameter κ , the wave functions correspond to the characters of the compact simple Lie groups ($\kappa = 1$) [We 1925/26] or to zonal spherical functions on symmetric spaces ($\kappa = 1/2, 2, 4$) [Ha 1958], [He 1978]. At arbitrary values of κ , they provide an interpolation between these objects.

This class has many remarkable properties. Here we mention only one: the wave functions of such systems are a natural generalization of special functions (hypergeometric functions) to the case of several variables. The history of this problem and some results may be found in [OP 1983]. In [Pe 1998a], it was shown that the product of two wave functions is a finite linear combination of analogous functions, namely, of functions that appear in the corresponding Clebsch–Gordan series. In other words, this deformation (κ -deformation) does not change the Clebsch–Gordan series. For rank 1, we obtain the well-known cases of the Legendre, Gegenbauer and Jacobi polynomials, and the limiting cases of the Laguerre and Hermite polynomials (see for example [Vi 1968]). Some other cases were also considered in [He 1955], [Ja 1970], [Ko 1974], [Ja 1975], [Vr 1976], [Se 1977], [KS 1978], [Ma 1982], [Pr 1984], [Vr 1984], [HO 1987], [He 1987], [Op 1988a], [Op 1988b], [Ma 1988], [La 1989], [Op 1989], [St 1989], [KS 1995] and [CP 1997]. In [PRZ 1998], κ -deformed Clebsch–Gordan series was used in order to obtain the explicit expressions for the generalized Gegenbauer polynomials³ of type A_2 what gives the explicit solution of the three-body Calogero–Sutherland model. For special values of $\kappa = 1/2, 2, 4$, these formulae give the explicit expressions for zonal polynomials of type A_2 .

In [Pe 1999] and Appendix B, are presented analogous results obtained by E. Ragoucy, Ph. Zaugg and the author for two-parameter family of polynomials of type A_2 introduced by Ruijsenaars [Ru 1987] and Macdonald [Ma 1988].

³ In some papers, the name Jack polynomials [Ja 1970] is used. However, the Jack polynomials are a very special case of polynomials under consideration. So, we prefer to use the name generalized Gegenbauer polynomials for the general case and Jack polynomials for the special case.

2 General description

The systems under consideration are described by the Hamiltonian (for more details, see [OP 1983])

$$H = \frac{1}{2} p^2 + U(q), \quad p^2 = (p, p) = \sum_{j=1}^l p_j^2, \quad (2.1)$$

where $p = (p_1, \dots, p_l)$, $p_j = -i \partial / \partial q_j$, is a momentum operator, and $q = (q_1, \dots, q_l)$ is a coordinate vector in the l -dimensional vector space $V \sim \mathbb{R}^l$ with standard scalar product $(\cdot, \cdot)$. The potential $U(q)$ is constructed by means of a certain system of vectors $R^+ = \{\alpha\}$ in V (the so-called *root systems*):

$$U = \sum_{\alpha \in R^+} g_\alpha^2 v(q_\alpha), \quad q_\alpha = (\alpha, q), \quad g_\alpha^2 = \kappa_\alpha(\kappa_\alpha - 1), \quad g_\alpha = g_\beta, \quad \text{if } (\alpha, \alpha) = (\beta, \beta). \quad (2.2)$$

Such systems are completely integrable for potentials of five types (see [OP 1983] for A_l ; [HO 1987], [He 1987], [Op 1988a], [Op 1988b] and [Op 1989] for a general case). They are a generalization of the Calogero–Sutherland systems [Ca 1971], [Su 1972] for which $\{\alpha\} = \{e_i - e_j\}$, $\{e_j\}$ being a standard basis in V .

In this paper, we consider in details only the case of A_2 with potential $v(q) = \sin^{-2} q$. For the description of other cases see [Pe 1999].

3 The Clebsch–Gordan series

Let us recall the main results of [Pe 1998a] and specialize them to the A_2 case with potential $v(q) = \sin^{-2} q$.

The Schrödinger equation for this quantum system has the form

$$H \Psi^\kappa = E(\kappa) \Psi^\kappa; \quad H = -\Delta_2 + U(q_1, q_2, q_3), \quad \Delta_2 = \sum_{j=1}^3 \frac{\partial^2}{\partial q_j^2} \quad (3.1)$$

with potential

$$U(q_1, q_2, q_3) = \kappa(\kappa - 1) \left(\sin^{-2}(q_1 - q_2) + \sin^{-2}(q_2 - q_3) + \sin^{-2}(q_3 - q_1) \right). \quad (3.2)$$

The ground state wave function and its energy are

$$\Psi_0^\kappa(q) = \left(\prod_{j < k}^3 \sin(q_j - q_k) \right)^\kappa, \quad E_0(\kappa) = 8 \kappa^2. \quad (3.3)$$

Substituting $\Psi_\lambda^\kappa = \Phi_\lambda^\kappa \Psi_0^\kappa$ in (3.1), we obtain

$$-\Delta^\kappa \Phi_\lambda^\kappa = \varepsilon_\lambda(\kappa) \Phi_\lambda^\kappa, \quad \Delta^\kappa = \Delta_2 + \Delta_1^\kappa, \quad \varepsilon_\lambda(\kappa) = E_\lambda(\kappa) - E_0(\kappa). \quad (3.4)$$

Here the operator Δ_1^κ takes the form

$$\Delta_1^\kappa = \kappa \sum_{j < k}^3 \cot(q_j - q_k) \left(\frac{\partial}{\partial q_j} - \frac{\partial}{\partial q_k} \right). \quad (3.5)$$

It is easy to see that the set of symmetric polynomials in variables $\exp(2iq_j)$ is invariant under the action of Δ^κ . Such polynomial m_λ is labelled by the $SU(3)$ highest weight $\lambda = m\lambda_1 + n\lambda_2$, with m, n being non-negative integers, and $\lambda_{1,2}$ being two fundamental weights. In general,

$$\Phi_\lambda^\kappa = \sum_{\mu \leq \lambda} C_\lambda^\mu(\kappa) m_\mu, \quad \mu, \lambda \in P^+, \quad m_\mu = \sum_{\nu \in W \cdot \mu} e^{2i(q, \nu)}, \quad (3.6)$$

where P^+ denotes the cone of dominant weights, W is the Weyl group, and $C_\lambda^\mu(\kappa)$ are some constants.

As it was shown in [Pe 1998a], the product of two wave functions is a finite sum of wave functions (a sort of the κ -deformed Clebsch–Gordan series):

$$\Phi_\mu^\kappa \Phi_\lambda^\kappa = \sum_{\nu \in D_\mu(\lambda)} C_{\mu\lambda}^\nu(\kappa) \Phi_\nu^\kappa. \quad (3.7)$$

In this equation, $D_\mu(\lambda) = (D_\mu + \lambda) \cap P^+$, where D_μ is a weight diagram of the representation with the highest weight μ .

Since Φ_μ^κ are symmetric functions of $\exp(2iq_j)$, it is convenient to use a new set of variables:

$$\begin{aligned} z_1 &= e^{2iq_1} + e^{2iq_2} + e^{2iq_3}, \\ z_2 &= e^{2i(q_1+q_2)} + e^{2i(q_2+q_3)} + e^{2i(q_3+q_1)}, \\ z_3 &= e^{2i(q_1+q_2+q_3)}. \end{aligned} \quad (3.8)$$

In the centre-of-mass frame ($\sum_i q_i = 0$), the wave functions depend only on two variables chosen as z_1 and z_2 (in this case, $z_3 = 1$). In these variables, up to a normalization factor, we have

$$\Delta^\kappa = (z_1^2 - 3z_2) \partial_1^2 + (z_2^2 - 3z_1) \partial_2^2 + (z_1 z_2 - 9) \partial_1 \partial_2 + (3\kappa + 1) (z_1 \partial_1 + z_2 \partial_2), \quad (3.9)$$

where $\partial_i = \partial / \partial z_i$. Corresponding eigenvalues are

$$\varepsilon_{m,n}(\kappa) = m^2 + n^2 + mn + 3\kappa(m + n). \quad (3.10)$$

We shall use the normalization for polynomials Φ_λ^κ such that the coefficient at the highest monomial is equal to one. Denoting them by $P_{m,n}^\kappa$, we have

$$P_{m,n}^\kappa(z_1, z_2) = \sum_{p,q} C_{m,n}^{p,q}(\kappa) z_1^p z_2^q = z_1^m z_2^n + \text{lower terms}, \quad (3.11)$$

with $p + q \geq m + n$ and $p - q \equiv m - n \pmod{3}$. As it is easy to see, the first polynomials are

$$P_{0,0}^\kappa = 1, \quad P_{1,0}^\kappa = z_1, \quad P_{0,1}^\kappa = z_2. \quad (3.12)$$

Simple consequences of (3.7) for $P_\lambda^\kappa = P_{1,0}^\kappa$ or $P_{0,1}^\kappa$ are [Pe 1998]

$$\begin{aligned} z_1 P_{m,n}^\kappa &= P_{m+1,n}^\kappa + a_{m,n}(\kappa) P_{m,n-1}^\kappa + c_m(\kappa) P_{m-1,n+1}^\kappa, \\ z_2 P_{m,n}^\kappa &= P_{m,n+1}^\kappa + \tilde{a}_{m,n}(\kappa) P_{m-1,n}^\kappa + c_n(\kappa) P_{m+1,n-1}^\kappa, \end{aligned} \quad (3.13)$$

where

$$\begin{aligned} a_{m,n}(\kappa) &= \tilde{a}_{n,m}(\kappa) = c_n(\kappa) c_{m+n+\kappa}(\kappa), \\ c_m(\kappa) &= \frac{e(m)}{e(\kappa + m)}, \quad e(m) = \frac{m}{m - 1 + \kappa}. \end{aligned} \quad (3.14)$$

Below we shall construct such polynomials using these recurrence relations.

4 A_2 case

Now we proceed to the case of $A_2 \sim su(3)$. In this case, the representation d of A_2 is characterized by two non-negative numbers $d = d_{mn}$. We have two fundamental representations

$$d_{10}, \quad d_{01}; \quad (\dim d_{10} = \dim d_{01} = 3),$$

and also representations

$$\begin{aligned} d_{n0}, \quad d_{0n}; \quad &\left(\dim d_{n0} = \dim d_{0n} = \frac{1}{2}(n+1)(n+2) \right), \\ d_{n1} \left(\dim d_{n1} = (n+1)(n+3) \right), \quad &d_{nn} \left(\dim d_{nn} = (n+1)^3 \right). \end{aligned}$$

We start with the Clebsch–Gordan series

$$\begin{aligned} d_{10} \otimes d_{n+1,0} &= d_{n+2,0} \oplus d_{n,1}, \\ d_{01} \otimes d_{n0} &= d_{n,1} \oplus d_{n-1,0}. \end{aligned} \quad (4.1)$$

Excluding $d_{n,1}$, we obtain

$$d_{n-1,0} \ominus (d_{01} \otimes d_{n0}) \oplus (d_{10} \otimes d_{n+1,0}) \ominus d_{n+2,0} = 0,$$

or

$$\chi_{n-1,0} - z_2 \chi_{n,0} + z_1 \chi_{n+1,0} - \chi_{n+2,0} = 0, \quad (4.2)$$

where are introduced the notations:

$$\begin{aligned} z_1 &= \chi_{10} = e^{i\theta_1} + e^{i\theta_2} + e^{i\theta_3}, \\ z_2 &= \chi_{01} = e^{-i\theta_1} + e^{-i\theta_2} + e^{-i\theta_3}. \end{aligned} \quad (4.3)$$

From this, we obtain the expression for the generating function

$$\begin{aligned} F_0^1(z_1, z_2; u) &= \sum_{n=0}^{\infty} \chi_{n0}(z_1, z_2) u^n, \\ F_0^1(z_1, z_2; u) &= (1 - z_1 u + z_2 u^2 - u^3)^{-1}. \end{aligned} \quad (4.4)$$

Let us define now the κ -deformed functions $\tilde{P}_{n,0}^\kappa(z_1, z_2)$ by the formula

$$F^\kappa(z_1, z_2; u) = (1 - z_1 u + z_2 u^2 - u^3)^{-\kappa} = \sum_{n=0}^{\infty} \tilde{P}_{n,0}^\kappa(z_1, z_2) u^n. \quad (4.5)$$

Differentiating F^κ on u , z_1 and z_2 , we get

$$\begin{aligned} F_u^\kappa &= \kappa (z_1 - 2z_2 u + 3u^2) F^{\kappa+1}, \\ F_{z_1, z_1}^\kappa &= \kappa(\kappa + 1) u^2 F^{\kappa+2}, & F_{z_2, z_2}^\kappa &= \kappa(\kappa + 1) u^4 F^{\kappa+2}, \\ F_{z_1, z_2}^\kappa &= -\kappa(\kappa + 1) u^3 F^{\kappa+2}, & u F_u^\kappa &= \kappa (z_1 u - 2z_2 u^2 + 3u^3) F^{\kappa+1}, \end{aligned}$$

and

$$(1 - z_1 u + z_2 u^2 - u^3) F_u^\kappa = \kappa (z_1 - 2z_2 u + 3u^2) F^\kappa. \quad (4.6)$$

From this it follows the important recurrence formula

$$(n+3) \tilde{P}_{n+3,0}^\kappa = (n+2+\kappa) z_1 \tilde{P}_{n+2,0}^\kappa - (n+1+2\kappa) z_2 \tilde{P}_{n+1,0}^\kappa + (n+3\kappa) \tilde{P}_{n,0}^\kappa. \quad (4.7)$$

We have also

$$F_{z_1}^\kappa = \kappa u F^{\kappa+1}, F_{z_2}^\kappa = -\kappa u^2 F^{\kappa+1}. \quad (4.8)$$

Hence

$$\partial_{z_1} P_{n,0}^\kappa = n P_{n-1,0}^{\kappa+1}, \partial_{z_2} P_{n,0}^\kappa = -\frac{n(n-1)}{\kappa+n-1} P_{n-2,0}^{\kappa+1}. \quad (4.9)$$

Finally we have the basic differential equation for $F^\kappa(z_1, z_2; u)$:

$$\begin{aligned} &((D_{z_1}^2 + D_{z_2}^2 + D_{z_1} D_{z_2}) - 3z_2 \partial_{z_1}^2 - 3z_1 \partial_{z_2}^2 - 9\partial_{z_1} \partial_{z_2} + 3\kappa(D_{z_1} + D_{z_2})) F^\kappa \\ &= (D_u^2 + 3\kappa D_u) F^\kappa(z_1, z_2; u); \end{aligned}$$

$$D_{z_1} = z_1 \partial_{z_1}, \quad D_{z_2} = z_2 \partial_{z_2}, \quad D_u = u \partial_u.$$

Let us note that the normalization of polynomials $\tilde{P}_{n,0}^\kappa(z_1, z_2)$ follows from the expression (4.5) for the generating function. Namely,

$$\tilde{P}_{n0}^\kappa(z_1, z_2) = \frac{(\kappa)_n}{n!} z_1^n + \dots = \frac{(\kappa)_n}{n!} P_{n,0}^\kappa(z_1, z_2), \quad (4.10)$$

where

$$(\kappa)_n = (\kappa)(\kappa + 1) \cdots (\kappa + n - 1).$$

The main property of this normalization is that $\tilde{P}_{n,0}^\kappa(z_1, z_2)$ has a polynomial dependence on the parameter κ .

Now we shall consider other Clebsch–Gordan series for $\kappa = 1$:

$$d_{1,0} \otimes d_{n+1,0} = d_{n+2,0} \oplus d_{n,1}.$$

According to [Pe 1998], the analogous formula is valid for an arbitrary value of κ , i.e.,

$$a_n z_1 \tilde{P}_{n+1,0}^\kappa = b_n \tilde{P}_{n+2,0}^\kappa + c_n \tilde{P}_{n,1}^\kappa. \quad (4.11)$$

Here

$$a_n = \kappa + n + 1, \quad b_n = n + 2,$$

and we have

$$c_n(\kappa) \tilde{P}_{n,1}^\kappa = (\kappa + n + 1) z_1 \tilde{P}_{n+1,0}^\kappa - (n + 2) \tilde{P}_{n+2,0}^\kappa. \quad (4.12)$$

Let us calculate now the generating function for both left-hand and right-hand sides of this equation,

$$G^\kappa = \sum_{n=0}^{\infty} c_n(\kappa) \tilde{P}_{n,1}^\kappa(z_1, z_2), \quad (4.13)$$

$$G^\kappa = \frac{\kappa z_1}{u} (F_0^\kappa - 1) + \frac{z_1}{u} F_1^\kappa - \frac{1}{u^2} (F_1^\kappa - \kappa z_1 u),$$

where

$$F_0^\kappa = \sum_{n=0}^{\infty} \tilde{P}_{n,0}^\kappa u^n; \quad F_1^\kappa = \sum_{n=0}^{\infty} n \tilde{P}_{n,0}^\kappa u^n = D_u F_0^\kappa,$$

and

$$G^\kappa = \frac{1}{u^2} (\kappa z_1 u - (1 - z_1 u) D_u) F_0^\kappa.$$

Finally,

$$G^\kappa = \kappa (2z_2 - (z_1 z_2 + 3)u + 2z_1 u^2) F_0^{\kappa+1}. \quad (4.14)$$

From this, it follows three-term recurrence relation

$$\tilde{P}_{n,1}^\kappa = \kappa \left(2z_2 \tilde{P}_{n,0}^{\kappa+1} - (z_1 z_2 + 3) \tilde{P}_{n-1,0}^{\kappa+1} + 2z_1 \tilde{P}_{n-2,0}^{\kappa+1} \right).$$

Now let us follow [PRZ 1998] and construct the general polynomials in terms of the simplest polynomials (Jack polynomials) $P_{m,0}^\kappa$ and $P_{0,n}^\kappa$. We get

$$P_{m,0}^\kappa P_{0,n}^\kappa = \sum_{i=0}^{\min(m,n)} \gamma_{m,n}^i P_{m-i,n-i}^\kappa. \quad (4.15)$$

This is a consequence of equation (3.7), with the notable difference that the sum on the right-hand side is over a restricted domain (actually, it parallels exactly the $SU(3)$ Clebsch–Gordan decomposition).

To prove this, let us assume that (4.15) is valid up to (m, n) . Then using (3.13) and (3.14), we get

$$P_{m,0}^\kappa P_{0,n+1}^\kappa = \sum_{i=0}^{\min(m,n+1)} \gamma_{m,n+1}^i P_{m-i,n+1-i}^\kappa + c_n \delta_{m,n+1}^i P_{m+1-i,n-1-i}^\kappa, \quad (4.16)$$

where we defined

$$\gamma_{m,n+1}^i = \gamma_{m,n}^i + \tilde{a}_{m-i+1,n-i+1} \gamma_{m,n}^{i-1} - c_n c_{m-i+1} \gamma_{m,n-1}^{i-1}, \quad (4.17)$$

$$\begin{aligned} \delta_{m,n+1}^i &= c_n^{-1} c_{n-i} \gamma_{m,n}^i - \gamma_{m,n-1}^i - a_{m-i+1,n-i} \gamma_{m,n-1}^{i-1} \\ &\quad + c_{n-1} c_{\kappa+n-1} \gamma_{m,n-2}^{i-1}. \end{aligned} \quad (4.18)$$

From the polynomial normalization, we already know that $\gamma_{m,n}^0 = 1$. After a straightforward computation, the solution to (4.17) proved to be equal

$$\gamma_{m,n}^i = \frac{e(2\kappa + m + n + 1 - i)_{-i}}{e(1)_i e(m+1)_{-i} e(n+1)_{-i}}, \quad (4.19)$$

where

$$e(m) = \frac{m}{m-1+\kappa}.$$

It implies that $\delta_{m,n+1}^i = 0$ in (4.18). Let us give also the more explicit expression⁴

$$\gamma_{m,n}^i = \frac{(\kappa)^i (m)_i (n)_i (3\kappa + m + n - 1 - i)_i}{i! (\kappa + m - 1)_i (\kappa + n - 1)_i (2\kappa + m + n - i)_i}, \quad (4.20)$$

where

$$\begin{aligned} (x)^i &= x(x+1) \cdots (x+i-1), \\ (x)_i &= x(x-1) \cdots (x-i+1). \end{aligned} \quad (4.21)$$

The constructive aspect of this formula is in its inverted form.

Theorem 1 [PRZ 1998]. *The generalized Gegenbauer polynomials $P_{m,n}^\kappa$ of type A_2 are given by the formula*

$$P_{m,n}^\kappa = \sum_{i=0}^{\min(m,n)} \beta_{m,n}^i P_{m-i,0}^\kappa P_{0,n-i}^\kappa, \quad (4.22)$$

⁴ Note that this expression for $\gamma_{m,n}^i$ may be obtained from the general Macdonald formula [Ma 1995], however, the way of proof given in [PRZ 1998] is more convenient here.

where the constants $\beta_{m,n}^i$ are

$$\beta_{m,n}^i = \frac{(-1)^i}{i!} \frac{3\kappa + m + n - 2i}{3\kappa + m + n - i} \frac{(m)_i (n)_i (\kappa)_i (3\kappa + m + n - 1)_i}{(\kappa + m - 1)_i (\kappa + n - 1)_i (2\kappa + m + n - 1)_i}. \quad (4.23)$$

Note that $\beta_{m,n}^i$ are obtained by using of the relation

$$\beta_{m,n}^i = - \sum_{j=0}^{i-1} \beta_{m,n}^j \gamma_{m-j,n-j}^{i-j}. \quad (4.24)$$

From this theorem, we see that the construction of a general polynomial $P_{m,n}^\kappa$ is similar to the construction of $SU(3)$ representations from tensor products of two fundamental representations.

Likewise, we can consider other types of decompositions, such as

$$P_{m,0}^\kappa P_{n,0}^\kappa = \sum_{i=0}^{\min(m,n)} \tilde{\gamma}_{m,n}^i P_{m+n-2i,i}^\kappa. \quad (4.25)$$

The proof is analogous to (4.15) (see footnote 3). The coefficients $\tilde{\gamma}_{m,n}^i$ are given by the formula

$$\tilde{\gamma}_{m,n}^i = \frac{(\kappa)^i (m)_i (n)_i (2\kappa + m + n - 1 - i)_i}{i! (\kappa + m - 1)_i (\kappa + n - 1)_i (\kappa + m + n - i)_i}. \quad (4.26)$$

Theorem 2 [PRZ 1998]. *There is another formula for polynomials $P_{m,n}^\kappa$ at $m \geq n$:*

$$\tilde{\gamma}_{m+n,n}^n P_{m,n}^\kappa = \sum_{i=0}^n \tilde{\beta}_{m,n}^i P_{m+n+i,0}^\kappa P_{n-i,0}^\kappa, \quad (4.27)$$

where

$$\tilde{\beta}_{m,n}^i = \frac{(-1)^i}{i!} (\kappa)_i \frac{(m+2i)}{m} \frac{(\kappa + m + n)^i}{(m+n+1)^i} \frac{(m)^i}{(\kappa + m + 1)^i} \frac{(n)_i}{(\kappa + n - 1)_i}. \quad (4.28)$$

This theorem follows directly from equation (4.25). The coefficients $\tilde{\beta}_{m,n}^i$ are found by using of the relation

$$\tilde{\beta}_{m,n}^i = - (\tilde{\gamma}_{m+n+i,n-i}^{n-i})^{-1} \sum_{j=0}^{i-1} \tilde{\beta}_{m,n}^j \tilde{\gamma}_{m+n+j,n-j}^{n-i}. \quad (4.29)$$

As a by-product, let us specialize equation (4.22) to the case $\kappa = 1$, where $P_{m,n}^\kappa$ are nothing but the $SU(3)$ characters. We get

$$P_{m,n}^1 = P_{m,0}^1 P_{0,n}^1 - P_{m-1,0}^1 P_{0,n-1}^1. \quad (4.30)$$

From this, we easily deduce the generating function for $SU(3)$ characters (see e.g. [PS 1978])

$$G^1(u, v) = \sum_{m,n=0}^{\infty} u^m v^n P_{m,n}^1 = \frac{1 - uv}{(1 - z_1 u + z_2 u^2 - u^3)(1 - z_2 v + z_1 v^2 - v^3)}. \quad (4.31)$$

Closing this section, let us note that for $\kappa=1/2, 1, 2$ and 4 , the obtained formulae yield the explicit expression of zonal polynomials for certain symmetric spaces listed below.

Let us mention also the papers [CP 1997], [Pe 1998b] where the integral representations for the case $N=3$ were obtained.

Appendix A. List of polynomials $P_{m,n}^\kappa$ with $m+n \leq 4$

Following [PRZ 1998], we list here the polynomials $P_{m,n}^\kappa$ with $m+n \leq 4$:

$$\begin{aligned} P_{2,0}^\kappa &= z_1^2 - \frac{2}{\kappa+1} z_2, \\ P_{1,1}^\kappa &= z_1 z_2 - \frac{3}{2\kappa+1}, \\ P_{3,0}^\kappa &= z_1^3 - \frac{6}{\kappa+2} z_1 z_2 + \frac{6}{(\kappa+1)(\kappa+2)}, \\ P_{2,1}^\kappa &= z_1^2 z_2 - \frac{2}{\kappa+1} z_2^2 - \frac{3\kappa+1}{(\kappa+1)^2} z_1, \\ P_{4,0}^\kappa &= z_1^4 - \frac{12}{\kappa+3} z_1^2 z_2 + \frac{12}{(\kappa+2)(\kappa+3)} z_2^2 + \frac{24}{(\kappa+2)(\kappa+3)} z_1, \\ P_{3,1}^\kappa &= z_1^3 z_2 - \frac{6}{\kappa+2} z_1 z_2^2 - \frac{3(3\kappa+2)}{(\kappa+2)(2\kappa+3)} z_1^2 + \frac{30}{(\kappa+2)(2\kappa+3)} z_2, \\ P_{2,2}^\kappa &= z_1^2 z_2^2 - \frac{2}{\kappa+1} (z_1^3 + z_2^3) - \frac{12(\kappa-1)}{(\kappa+1)(2\kappa+3)} z_1 z_2 + \frac{9(\kappa-1)}{(\kappa+1)^2(2\kappa+3)}. \end{aligned}$$

Appendix B. Some formulae for Macdonald polynomials for A_2 case (by A.M. Perelomov, E. Ragoucy and Ph. Zaugg)

Here we give only some necessary for us information. For other results and details, see [Ma 1995].

The Macdonald polynomials of type A_2 may be defined as polynomial eigenfunctions of the Macdonald difference equation

$$M^1 P_{m,n}^{(q,t)}(x_1, x_2, x_3) = \lambda P_{m,n}^{(q,t)}(x_1, x_2, x_3), \quad P_{m,n} = z_1^m z_2^n + \text{lower terms}, \quad (\text{B.1})$$

$$z_1 = x_1 + x_2 + x_3, \quad z_2 = x_1x_2 + x_2x_3 + x_3x_1, \quad z_3 = x_1x_2x_3,$$

where

$$M^1 = \prod_{j \neq k} \frac{(tx_j - x_k)}{(x_j - x_k)} T_j, \quad T_1 f(x_1, x_2, x_3) = f(qx_1, x_2, x_3), \dots \quad (\text{B.2})$$

The Macdonald polynomials satisfy the following recurrence relation

$$z_1 P_{m,n}^{(q,t)}(z_1, z_2, z_3) = P_{m+1,n}^{(q,t)} + a_{m,n}(q, t) P_{m,n-1}^{(q,t)} + b_{m,n}(q, t) P_{m-1,n+1}^{(q,t)}, \quad (\text{B.3})$$

where

$$\begin{aligned} a_{mn} &= c_n \tilde{c}_{m+n}, & b_{mn} &= c_m, \\ c_m(q, t) &= \frac{(1 - q^m)(1 - t^2 q^{m-1})}{(1 - tq^m)(1 - tq^{m-1})}, \\ \tilde{c}_{m+n}(q, t) &= \left(\frac{1 - tq^{m+n}}{1 - t^2 q^{m+n}} \right) \left(\frac{1 - t^3 q^{m+n-1}}{1 - t^2 q^{m+n-1}} \right). \end{aligned} \quad (\text{B.4})$$

Note that

$$c_m(q, 1) = c_m(q, q) = 1, \quad \tilde{c}_{m+n}(q, 1) = \tilde{c}_{m+n}(q, q) = 1. \quad (\text{B.5})$$

Now let us consider the case $t = q^k$, k being an integer. In this case, the explicit expression for the generating function of polynomials $P_{n,0}^{(k)} \equiv P_{n,0}^{(q,t)}$ has the form

$$\begin{aligned} G^{(k)}(u) &= \prod_{j=0}^{k-1} F^1(q^j u), \\ G^{(k)}(u) &= \sum C_n^k P_{n,0}^{(k)} u^n, \\ C_n^k &= \frac{[k]_n}{[1]_n}, \quad [x]_n = [x][x+1] \cdots [x+n-1], \quad [x] = \frac{1 - q^x}{1 - x}. \end{aligned} \quad (\text{B.6})$$

From this we get a three-term recurrence relation being analogous to (4.7)

$$[n+1] \tilde{P}_{n+1}^k = [k+n] z_1 \tilde{P}_n^k - [2k+n-1] z_2 \tilde{P}_{n-1}^k + [3k+n-2] z_3 \tilde{P}_{n-2}^k, \quad (\text{B.7})$$

which follows from the recurrence relation

$$(1 - z_1 u + z_2 u^2 - z_3 u^3) G^{(k)}(u) = (1 - z_1 u q^k + z_2 u^2 q^{2k} - z_3 u^3 q^{3k}) G(qu). \quad (\text{B.8})$$

From here, we obtain

$$\tilde{P}_0^\kappa = 1, \quad \tilde{P}_1^\kappa = [\kappa] z_1, \quad \tilde{P}_2^\kappa = \frac{[\kappa+1][\kappa]}{[1][2]} z_1^2 - \frac{[2\kappa]}{[2]} z_2, \dots \quad \tilde{P}_{m,0} = \frac{[1]_m}{[\kappa]_m} P_{m,0},$$

Let us give also the formula for the generating function for the case of arbitrary t :

$$G^{(q,t)}(u) = \prod_{j=0}^{\infty} \frac{(1 - q^j t u)}{(1 - q^j u)} = \sum_{j=0}^{\infty} c_j(q, t) u^j. \quad (\text{B.9})$$

Using the results from the book [Ma 1995], it is not difficult to get the formulae

$$P_{m,0}^{(q,t)} P_{0,n}^{(q,t)} = \sum_{i=0}^{\min(m,n)} \gamma_{m,n}^i P_{m-i,n-i}^{(q,t)}, \quad (\text{B.10})$$

where

$$\gamma_{m,n}^i = \frac{(t; q)_i (q^m; q^{-1})_i (q^n; q^{-1})_i (q^{m+n-1-i} t^3; q^{-1})_i}{(q; q)_i (q^{m-1} t; q^{-1})_i (q^{n-1} t; q^{-1})_i (q^{m+n-i} t^2; q^{-1})_i}; \quad (\text{B.11})$$

and

$$P_{m,0}^{(q,t)} P_{n,0}^{(q,t)} = \sum_{i=0}^{\min(m,n)} \tilde{\gamma}_{m,n}^i P_{m+n-2i,i}^{(q,t)}, \quad (\text{B.12})$$

where

$$\tilde{\gamma}_{m,n}^i = \frac{(t; q)_i (q^m; q^{-1})_i (q^n; q^{-1})_i (q^{m+n-i-1} t^2; q^{-1})_i}{(q; q)_i (q^{m-1} t; q^{-1})_i (q^{n-1} t; q^{-1})_i (q^{m+n-i} t; q^{-1})_i}. \quad (\text{B.13})$$

Inverting formulae (B.10) and (B.11) as in the [PRZ 1998], we come to

Theorem 1a. *The Macdonald polynomials $P_{m,n}^{(q,t)}$ of type A_2 are given by the formula*

$$P_{m,n}^{(q,t)} = \sum_{i=0}^{\min(m,n)} \beta_{m,n}^i P_{m-i,0}^{(q,t)} P_{0,n-i}^{(q,t)}, \quad (\text{B.14})$$

where constants have the form

$$\begin{aligned} \beta_{m,n}^i &= (-1)^i q^{i(i-1)/2} \frac{(t; q^{-1})_i}{(q; q)_i} \frac{1 - q^{m+n-2i} t^3}{1 - q^{m+n-i} t^3} \\ &\times \frac{(q^m; q^{-1})_i (q^n; q^{-1})_i (q^{m+n-1} t^3; q^{-1})_i}{(q^{m-1} t; q^{-1})_i (q^{n-1} t; q^{-1})_i (q^{m+n-1} t^2; q^{-1})_i}. \end{aligned}$$

In similar way, we get

Theorem 2a. *The Macdonald polynomials $P_{m,n}^{(q,t)}$ of type A_2 are given by the formula*

$$\tilde{\gamma}_{m+n,n}^n P_{m,n}^{(q,t)} = \sum_{i=0}^n \tilde{\beta}_{m,n}^i P_{m+n+i,0}^{(q,t)} P_{n-i,0}^{(q,t)}, \quad m \geq n \quad (\text{B.15})$$

where constants $\tilde{\gamma}_{m,n}^i$ are given by formula (B.13), and

$$\tilde{\beta}_{m,n}^i = (-1)^i q^{i(i-1)/2} \frac{(t; q^{-1})_i}{(q; q)_i} \frac{(1 - q^{m+2i})}{(1 - q^m)} \frac{(q^m; q)_i (q^n; q^{-1})_i (q^{m+n} t; q)_i}{(q^{m+n+1}; q)_i (q^{m+1} t; q)_i (q^{n-1} t; q^{-1})_i}. \quad (\text{B.16})$$

From Theorems 1a and 2a, many interesting identities may be obtained. Here we give one of them:

$$S_{n,l}^k = \sum_{i=0}^l (-1)^i q^{i(i-1)/2} \frac{[3k+n-2i]}{[3k+n-i]} \left(\prod_{j=1}^i \frac{[k-j+1]}{[j]} \frac{[3k+n-j]}{[2k+n-j]} \right) \\ \times \left(\prod_{j=0}^{l-i-1} \frac{[k+j]}{[j+1]} \frac{[3k+n-l-i-j-1]}{[2k+n-l-i-j]} \right) = 0, \quad (\text{B.17})$$

where $[n] = (1 - q^n)/(1 - q)$.

The simplest version of this formula is

$$\sum_{i=0}^l (-1)^i q^{i(i-1)/2} \frac{[k+l-i-1]}{[i]! [l-i]! [k-i]!} = 0. \quad (\text{B.18})$$

We give below the **list of polynomials** $P_{m,n}^{(q,t)}$ at $m+n \leq 4$

$$\begin{aligned} P_{0,0}^{(q,t)} &= 1, & P_{1,0}^{(q,t)} &= z_1, & P_{0,1}^{(q,t)} &= z_2, \\ P_{2,0}^{(q,t)} &= z_1^2 - \frac{(1-q)(1+t)}{(1-qt)} z_2, \\ P_{1,1}^{(q,t)} &= z_1 z_2 - \frac{(1-q)(1+t+t^2)}{(1-qt^2)} z_3, \\ P_{3,0}^{(q,t)} &= z_1^3 - \frac{(1-q)(2+q+t+2qt)}{1-q^2t} z_1 z_2 + \frac{(1-q)^2(1+q)(1+t+t^2)}{(1-qt)(1-q^2t)} z_3, \\ \\ P_{2,1}^{(q,t)} &= z_1^2 z_2 - \frac{(1-q)(1+t)}{1-qt} z_2^2 + \frac{(1-q^2)(1-qt^3)}{(1-qt)^2(1+qt)} z_1 z_3, \\ P_{4,0}^{(q,t)} &= z_1^4 - \frac{(1-q)(3+2q+q^2+t+2qt+3q^2t)}{1-q^3t} z_1^2 z_2 \\ &\quad + \frac{(1-q)^2(1+q+q^2)(1+t)(1+qt)}{(1-q^2t)(1-q^3t)} z_2^2 \\ &\quad + \frac{(1-q)^2(1+q)(2+q+q^2+t+2qt+q^2t+t^2+qt^2+2q^2t^2)}{(1-q^2t)(1-q^3t)} z_1 z_3, \\ \\ P_{3,1}^{(q,t)} &= z_1^3 z_2 - \frac{(1-q)(2+q+t+2qt)}{1-q^2t} z_1 z_2^2 - \frac{(1-q^3)(1-q^2t^3)}{(1-q^2t)(1-q^3t^2)} z_1^2 z_3 \\ &\quad + \frac{(1-q)^2}{(1-q^2t)(1-q^3t^2)} \\ &\quad \times (2+2q+q^2+2t+4qt+3q^2t+q^3t+t^2+ \\ &\quad 3qt^2+4q^2t^2+2q^3t^2+qt^3+2q^2t^3+2q^3t^3) z_2 z_3; \end{aligned}$$

$$\begin{aligned}
P_{2,2}^{(q,t)} &= z_1^2 z_2^2 - \frac{(1-q)(1+t)}{1-qt} (z_2^3 + z_1^3 z_3) \\
&\quad - \frac{(1-q)(3q+q^2-3t+2q^2t+q^3t-t^2-2qt^2+3q^3t^2-qt^3-3q^2t^3)}{(1-qt)(1-q^3t^2)} z_1 z_2 z_3 \\
&\quad + \frac{(1-q)^2(1+q)(1+t+t^2)(q-t+q^2t-qt^2+q^3t^2-q^2t^3)}{(1-qt)(1-q^2t^2)(1-q^3t^2)} z_3^2.
\end{aligned}$$

Acknowledgment

The paper was complete during my stay at Max-Planck-Institut für Mathematik Bonn. I am grateful the members of the Institute for hospitality.

References

- [CP 1997] Cariñena J.M. and Perelomov A.M., J. Phys. **A30**, L495–L501
- [Ca 1971] Calogero F., J. Math. Phys. **12**, 419–436
- [HO 1987] Heckman G. and Opdam E., Composito Math. **64**, 329–352
- [Ha 1958] Harish-Chandra, Amer. J. Math. **80**, 241–310 and 553–613
- [He 1955] Herz C., Ann. of Math. **61**, 474–523
- [He 1978] Helgason S., *Differential Geometry, Lie Groups and Symmetric Spaces*, Acad. Press: N.Y.
- [He 1987] Heckman G., Composito Math. **64**, 353–373
- [Hu 1972] Humphreys J.E., *Introduction to Lie Algebras and Representation Theory*, Springer
- [Ja 1970] Jack H., Proc. Roy. Soc. Edinb. **A69**, 1–18
- [Ja 1975] James A., pp 497–520 in: *Theory and Application of Special Functions*, Acad. Press: N.Y.
- [KS 1978] Koornwinder T. and Sprinkhuizen–Kuyper I., SIAM J. Math. Anal. **9**, 457–483
- [KS 1995] Kuznetsov V.B. and Sklyanin E.K., RIMS Kokyuroku **919**, 27–34
- [Ko 1974] Koornwinder T., Proc. Ned. Akad. Wet. **A77**, 48–66, 357–381
- [La 1989] Lassalle M., C.R. Acad. Sci., ser. I, **309**, 941–944

- [Ma 1982] Macdonald I.G., SIAM J.Math.Anal. **13**, 988–1007
- [Ma 1988] Macdonald I.G., *Orthogonal polynomials associated with root systems*, preprint
- [Ma 1995] Macdonald I.G., *Symmetric Functions and Hall Polynomilas*, Oxford Univ. Press
- [OP 1977] Olshanetsky M.A. and Perelomov A.M., Lett. Math. Phys. **2**, 7–13
- [OP 1978] Olshanetsky M.A. and Perelomov A.M., Funct. Anal. Appl. **12**, 121–128
- [OP 1983] Olshanetsky M.A. and Perelomov A.M., Phys. Rep. **94**, 313–404
- [OV 1990] Onishchik A.L. and Vinberg E.B., *Lie Groups and Algebraic Groups*, Springer–Verlag
- [Op 1988a] Opdam E., Composito Math. **67**, 21–49
- [Op 1988b] Opdam E., Composito Math. **67**, 191–209
- [Op 1989] Opdam E., Inv. Math. **98**, 1–18
- [PRZ 1998] Perelomov A.M., Ragoucy E. and Zaugg Ph., J. Phys. **A31**, L559–L565
- [PS 1978] Patera J. and Sharp R.T., Proceedings of the VII International Colloquium on Group Theoretical Methods in Physics, Austin, Texas.
- [Pe 1998a] Perelomov A.M., J. Phys. **A31**, L31–L37
- [Pe 1998b] Perelomov A.M., in: Proc. of the II International Workshop *Lie Theory and its Applications in Physics*. II, pp.171–183, Singapore: World Scientific
- [Pe 1999] Perelomov A.M., J. Phys. **A32**, 8563–8576
- [Pr 1984] Prati M.–C., Lett. Nuovo Cim. **41**, 275
- [Ru 1987] Ruijsenaars S., Comm. Math. Phys. **110**, 191–213
- [Se 1977] Sekiguchi J., Publ. RIMS Kyoto Univ. Suppl. **12**, 455–459
- [St 1989] Stanley R.P., Adv. Math. **77**, 76–115
- [Su 1972] Sutherland B., Phys. Rev. **A4**, 2019–2021
- [Vi 1968] Vilenkin N.Ja., *Special Functions and the Theory of Group Representations*, Amer. Math. Soc. Transl. Monographs **22**, Providence, RI

- [Vr 1976] Vretare L., Math. Scand. **39**, 343–358
- [Vr 1984] Vretare L., SIAM J. Math. Anal. **15**, 805–833
- [We 1925] Weyl H., Math. Zs. **23**, 271–309
- [We 1926] Weyl H., Math. Zs. **24**, 328–376 and 377–395